

ARECACEAE – PALM FAMILY

Plant: woody vines, shrubs and trees

Stem: branched or unbranched, woody

Root:

Leaves: 1 or 2 compound (fan-shaped or palmate) or feather shaped (pinnate) with a conspicuous stalk (petiole), sheathed at base (usually split at maturity), base of stalk often remains on trunk or stem

Flowers: perfect and/or imperfect (monoecious or dioecious), regular (actinomorphic); flowers small and whitish; 2-4 sepals, petals and few to many stamens; ovary superior, 1-3 pistils, 3 styles

Fruit: berry or leathery-like berry or drupe, any size with one large seed

Other: very large family, mostly in tropics to subtropics; Monocotyledons Group

Genera: 200+ genera; locally *Sabal* (palmetto)

ARECACEAE – PALM FAMILY

Cabbage Palmetto; *Sabal palmetto* (Walt.) Lodd. ex J.A. & J.H. Schultes

Cabbage Palmetto

Sabal palmetto (Walt.) Lodd.

ex J.A. & J.H. Schultes

Areaceae (Palm Family)

Fort Pulaski National Monument, Georgia

Notes: shrub or tree; flowers yellowish-white; a fan-leaved palm tree, leaf stalks continue into frond, at trunk end the leafstalk base is forked, loose fibers common on leaves, without spines on leaf stalk; fruit 0.5 in; state tree of Florida

[V Max Brown, 2006]

